

# PTGER1 and PTGER2 receptors mediate regulation of progesterone synthesis and type 1 11 $\beta$ -hydroxysteroid dehydrogenase activity by prostaglandin E<sub>2</sub> in human granulosa–lutein cells

C Chandras<sup>1,2</sup>, T E Harris<sup>1,2</sup>, A López Bernal<sup>3</sup>, D R E Abayasekara<sup>2</sup> and A E Michael<sup>1,4</sup>

<sup>1</sup>Department of Biochemistry and Molecular Biology, Royal Free and University College Medical School, University College London, Rowland Hill Street, London NW3 2PF, UK

<sup>2</sup>Department of Veterinary Basic Science, Royal Veterinary College, Royal College Street, London NW1 0TU, UK

<sup>3</sup>Department of Clinical Science at South Bristol (Obstetrics and Gynaecology), University of Bristol, Dorothy Hodgkin Building, Whitson Street, Bristol BS1 3NY, UK

<sup>4</sup>Division of Clinical Developmental Sciences, Academic Section of Obstetrics and Gynaecology, Centre for Developmental and Endocrine Signalling, St George's University of London, Cranmer Terrace, Tooting, London SW17 0RE UK

(Requests for offprints should be addressed to A E Michael; Email: [tony.michael@sgul.ac.uk](mailto:tony.michael@sgul.ac.uk))

(C Chandras is now at Developmental Biology, Foundation for Biomedical Research of the Academy of Athens, 11527 Athens, Greece)

## Abstract

In luteinizing granulosa cells, prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) can exert luteotrophic actions, apparently via the cAMP signalling pathway. In addition to stimulating progesterone synthesis, PGE<sub>2</sub> can also stimulate oxidation of the physiological glucocorticoid, cortisol, to its inactive metabolite, cortisone, by the type 1 11 $\beta$ -hydroxysteroid dehydrogenase (11 $\beta$ HSD1) enzyme in human granulosa–lutein cells. Having previously shown these human ovarian cells to express functional G-protein coupled, E-series prostaglandin (PTGER)1, PTGER2 and PTGER4 receptors, the aim of this study was to delineate the roles of PTGER1 and PTGER2 receptors in mediating the effects of PGE<sub>2</sub> on steroidogenesis and cortisol metabolism in human granulosa–lutein cells. PGE<sub>2</sub>-stimulated concentration-dependent increases in both progesterone production and cAMP accumulation (by  $1.9 \pm 0.1$ - and  $18.7 \pm 6.8$ -fold respectively at 3000 nM PGE<sub>2</sub>). While a selective PTGER1 antagonist, SC19220, could partially inhibit the steroidogenic response to PGE<sub>2</sub> (by

$55.9 \pm 4.1\%$  at 1000 nM PGE<sub>2</sub>), co-treatment with AH6809, a mixed PTGER1/PTGER2 receptor antagonist, completely abolished the stimulation of progesterone synthesis at all tested concentrations of PGE<sub>2</sub> and suppressed the stimulation of cAMP accumulation. Both PGE<sub>2</sub> and butaprost (a preferential PTGER2 receptor agonist) stimulated concentration-dependent increases in cortisol oxidation by 11 $\beta$ HSD1 (by  $42.5 \pm 3.1$  and  $40.0 \pm 3.0\%$  respectively, at PGE<sub>2</sub> and butaprost concentrations of 1000 nM). Co-treatment with SC19220 enhanced the ability of both PGE<sub>2</sub> and butaprost to stimulate 11 $\beta$ HSD1 activity (by  $30.2 \pm 0.2$  and  $30.5 \pm 0.6\%$  respectively), whereas co-treatment with AH6809 completely abolished the 11 $\beta$ HSD1 responses to PGE<sub>2</sub> and butaprost. These findings implicate the PTGER2 receptor–cAMP signalling pathway in the stimulation of progesterone production and 11 $\beta$ HSD1 activity by PGE<sub>2</sub> in human granulosa–lutein cells.

*Journal of Endocrinology* (2007) **194**, 595–602

## Introduction

Prostaglandins (PGs) have been implicated in the co-ordination of ovarian function, particularly in ovulation and in regulating the functional lifespan of the corpus luteum. For example, PGF<sub>2 $\alpha$</sub>  has luteolytic effects and can trigger functional luteal regression in a range of species, including humans and non-human primates (reviewed by Michael *et al.* 1994, Olofsson & Leung 1994), whereas selected PGs, like PGE<sub>2</sub>, can exert luteotrophic actions in human and non-human primate ovarian cells, stimulating progesterone synthesis apparently via the cAMP signalling pathway (Richardson 1986, Michael *et al.* 1993, 1994, Olofsson &

Leung 1994). Although PGs are established as major paracrine agents in ovarian physiology, their exact modes of action still remain to be elucidated.

Glucocorticoid steroids are also known to be important for gonadal function and have been implicated in the control of oocyte maturation (Greeley *et al.* 1986, Harlow *et al.* 1997, Yang *et al.* 1999, Chen *et al.* 2000). In potential target tissues, the physiological glucocorticoids, cortisol and corticosterone, are oxidised to inert metabolites (cortisone and 11-dehydrocorticosterone respectively) by 11 $\beta$ -hydroxysteroid dehydrogenase (11 $\beta$ HSD) enzymes (Seckl & Walker 2001, Seckl 2004, Draper & Stewart 2005). These enzymes are expressed and appear to play significant physiological roles in

reproductive tissues (reviewed by Michael *et al.* (2003)). For example, in the uterine-placental complex, both the relatively low-affinity NADP(H)-dependent type 1 11 $\beta$ HSD enzyme (11 $\beta$ HSD1) and the high-affinity NAD<sup>+</sup>-dependent type 2 11 $\beta$ HSD enzyme (11 $\beta$ HSD2) have recently been implicated in the physiological mechanism of labour (Challis *et al.* 2000, Whittle *et al.* 2001), although prior studies of human placenta and decidua had found no significant changes in cortisol metabolism by 11 $\beta$ HSD at parturition (López Bernal *et al.* 1982a,b).

Previously, we have established that luteinizing human granulosa cells, recovered from the ovarian follicular aspirates of women undergoing oocyte retrieval for *in vitro* fertilisation, exclusively express 11 $\beta$ HSD1 with no detectable expression of 11 $\beta$ HSD2 at either the mRNA or the protein level (Michael *et al.* 1997, Thurston *et al.* 2003a). Although 11 $\beta$ HSD1 generally acts as a 11-ketosteroid reductase to regenerate cortisol from cortisone (Seckl & Walker 2001, Seckl 2004, Draper & Stewart 2005), we have previously reported that in intact human granulosa-lutein cells in primary culture, 11 $\beta$ HSD1 acts predominantly as a 11 $\beta$ -dehydrogenase enzyme to catalyse the oxidative inactivation of cortisol (Michael *et al.* 1997, Thurston *et al.* 2003a). Since the utero-placental feedback loop for the induction of parturition (at least in sheep) appears to require stimulation of 11 $\beta$ HSD1 by PGs (Alfaidy *et al.* 2001, Challis *et al.* 1997), we have investigated the possible role for local PGs in control of 11 $\beta$ HSD1 in human granulosa-lutein cells *in vitro*. We recently reported that inhibition of local PG synthesis (using four structurally distinct inhibitors of PGH synthase) could suppress cortisol oxidation by 11 $\beta$ HSD1 in these human cells, but that this enzyme activity could be stimulated by PGE<sub>2</sub> (Jonas *et al.* 2006).

At the cellular level, PGE<sub>2</sub> actions are mediated through G protein-coupled, E-series PG (PTGER) receptors. To date, four subtypes of PTGER receptors have been cloned, designated PTGER1, PTEGER2, PTGER3 and PTGER4 (Hirata *et al.* 1991, Sugimoto *et al.* 1992, Funk *et al.* 1993, Honda *et al.* 1993, Watabe *et al.* 1993), with multiple isoforms of the PTGER3 receptor subtype (reviewed by Coleman *et al.* (1994)). While PTGER1 and PTGER3 receptors mediate increases in intracellular calcium concentrations, PTGER2 and PTGER4 receptors mediate stimulation of intracellular cAMP accumulation (Narumiya *et al.* 1999). Since PTGER receptors have different affinities for synthetic PGE<sub>2</sub> analogues, agonists and antagonists with differing degrees of preference or selectivity for each PTGER receptor subtype can be used to delineate the role for particular PTGER receptors in a given cellular response (Coleman *et al.* 1994). Using RT-PCR and preferential receptor agonists/antagonists, we and other researchers have established that human granulosa-lutein cells express PTGER1, PTGER2 and PTGER4 receptors, and that the PTGER1 and PTGER2 receptors are able to elicit increases in intracellular calcium and cAMP respectively (Harris *et al.* 2001, Narko *et al.* 2001). However, the exact PG receptors through which PGE<sub>2</sub> exerts its effects on progesterone synthesis and cortisol-cortisone

interconversion by 11 $\beta$ HSD1 have not been previously examined in ovarian cells. Hence, the aim of the current study was to examine the potential roles for PTGER1 and PTGER2 receptors in mediating the effects of PGE<sub>2</sub> on steroidogenesis and cortisol metabolism using human granulosa-lutein cells as our *in vitro* ovarian model.

## Materials and Methods

### *Isolation and culture of human granulosa cells*

Human granulosa cells were isolated from follicular aspirates of women undergoing controlled ovarian hyperstimulation for assisted conception at the Lister Private Hospital (Chelsea, London, UK) as previously described by Webley *et al.* (1988). Follicular aspirates were collected with informed patient consent in accordance with the Declaration of Helsinki and as approved by the local ethics committee.

Following isolation on 60% (v/v) Percoll, human granulosa-lutein cells were cultured for 48 h in 1:1 Dulbecco's modified Eagle's medium (DMEM):Ham's F12 medium supplemented with 10% (v/v) fetal calf serum (Invitrogen Life Technologies), 2 mM L-glutamine (Life Technologies), penicillin (87 000 IU/l; Sigma-Aldrich) and streptomycin (87 mg/l; Sigma-Aldrich) in an atmosphere of 5% (v/v) CO<sub>2</sub> in air. Cells were cultured either in 250  $\mu$ l volumes at a density of  $1 \times 10^5$  viable cells/well for assessing regulation of progesterone production and cAMP accumulation or in 1 ml volumes at a density of  $5 \times 10^4$  viable cells/well in 24-well plates for measurement of 11 $\beta$ HSD1 activities. For all experiments, cell viability, assessed by exclusion of 0.4% (v/v) trypan blue dye, was consistently > 85%.

### *Concentration-dependent effects of PGE<sub>2</sub> on progesterone production and cAMP accumulation in the absence or presence of SC19220 and AH6809*

In order to examine whether PTGER2 receptors were involved in mediating the steroidogenic response to PGE<sub>2</sub>, cells were transferred to serum-free 1:1 DMEM:Ham's F<sub>12</sub> medium supplemented with 1  $\mu$ M meclofenamic acid (Sigma-Aldrich) to suppress intrinsic PG synthesis (Jonas *et al.* 2006). Cells were challenged for 24 h with a range of PGE<sub>2</sub> concentrations (0–3000 nM; Cayman Chemical Company, MI, USA) in the absence and presence of either 10  $\mu$ M SC19220 or 10  $\mu$ M AH6809 (Alexis Biochemicals, Nottingham, UK). SC19220 and AH6809 were prepared to a stock concentration of 10 mM in dimethylsulphoxide (DMSO), the final concentration of which was diluted in all wells to 0.1% by volume.

After the 24-h treatment period, culture plates were frozen at  $-20^\circ\text{C}$ . Samples were subsequently thawed and progesterone and cAMP concentrations were measured using RIAs previously described by Pallikaros *et al.* (1995) and Steiner *et al.* (1972) respectively. The progesterone RIA had a working range of 0.5–8.0 nM with inter- and intra-assay coefficients of variation of 9 and 14% respectively. The cAMP

RIA included the acetylation step proposed by Harper & Brooker (1975), which gave the assay a working range of 0.2–1.3 nM with inter- and intra-assay coefficients of variation of 9 and 5% respectively.

*Effects of SC19220 and AH6809 on 11 $\beta$ HSD1 activities in response to PGE<sub>2</sub> and butaprost in human granulosa–lutein cells*

As in the assessment of the effects on progesterone production and cAMP accumulation, cells were pre-incubated for 48 h in serum-supplemented 1:1 DMEM:Ham's F12 medium. Following this, the cells were transferred to serum-free 1:1 DMEM:Ham's F12 medium, supplemented with 1  $\mu$ M meclofenamic acid and containing PGE<sub>2</sub> or butaprost (Cayman Chemical Company; 0–1000 nM of each), each in the absence or presence of 10  $\mu$ M SC19220 or AH6809 for 4 h. To overcome potential confounding effects of progesterone on 11 $\beta$ HSD activities (Thurston *et al.* 2003b), these experiments were performed in the presence of 100  $\mu$ M aminoglutethimide, having confirmed that this effectively suppresses progesterone output from human granulosa–lutein cells (Fowkes *et al.* 2001) without affecting cAMP accumulation. During the 4 h over which cells were exposed to PGE<sub>2</sub>/butaprost  $\pm$  SC19220/AH6809, 11 $\beta$ HSD1 activities were assessed using the radiometric conversion assay as previously described by Michael *et al.* (1995). In brief, intact cells in primary culture were co-incubated with 100 nM [1,2,6,7-<sup>3</sup>H]-cortisol (Amersham Biosciences), diluted to a specific activity of 0.5  $\mu$ Ci/100 pmol using non-radiolabelled cortisol (Sigma–Aldrich). At the end of the 4-h treatment period, spent culture medium was transferred to screw cap borosilicate tubes in which steroids were extracted by the addition of two volumes of chloroform (Merck). After evaporation to dryness under nitrogen gas at 45 °C, the steroid residues were suspended in 25  $\mu$ l ethyl acetate containing 1 mM cortisol and 1 mM cortisone (Sigma–Aldrich) before being resolved by thin layer chromatography (TLC) on silica 60 TLC plates (Merck), developed in an atmosphere of 92:8 chloroform:95% (v/v) ethanol. [<sup>3</sup>H]-cortisol and [<sup>3</sup>H]-cortisone were quantified using a Bioscan 200 TLC radiochromatogramme scanner (Lablogic, Sheffield, UK) and the oxidative activities of 11 $\beta$ HSD1 were calculated as net pmoles cortisol oxidised to cortisone over 4 h (Michael *et al.* 1995).

*Statistical analyses*

All experimental data are presented as the mean  $\pm$  S.E.M. for up to six independent experiments, where each experimental replicate was performed using cells from individual patients and each experimental condition was repeated either in quadruplicate or in triplicate within an experiment. Due to differences in absolute levels of progesterone production, cAMP accumulation and enzyme activities between patients, the results for each experiment were standardised and

presented as a percentage of basal control. However, all statistical analyses were performed using absolute data.

In all experiments, a one-way ANOVA with repeated measures was performed, followed by Bonferroni's multiple comparison as the *post hoc* test. All statistical evaluations were performed using GraphPad Prism version 3.02 (GraphPad Software Inc., San Diego, CA, USA) and significance was assessed in all experiments as a probability value of  $P < 0.05$ .

## Results

*Effects of SC19220 and AH6809 on progesterone production and 11 $\beta$ HSD1 activity*

Before use, both PTGER receptor antagonists were tested for possible direct effects on basal progesterone production and 11 $\beta$ HSD1 activity in human granulosa–lutein cells. Under conditions in which intrinsic PG synthesis had been suppressed with 1  $\mu$ M meclofenamic acid, incubation for 24 h with the preferential PTGER1 receptor antagonist SC19220 (10  $\mu$ M) and the PTGER1/PTGER2 receptor antagonist AH6809 (10  $\mu$ M) had no significant effect on basal progesterone production (over 24 h) or the oxidative activity of 11 $\beta$ HSD1 (over 4 h; Table 1).


*Concentration-dependent effects of PGE<sub>2</sub> on progesterone production in the absence or presence of SC19220 and AH6809*

Treatment with PGE<sub>2</sub> for 24 h resulted in a concentration-dependent increase in progesterone synthesis (Fig. 1). At PGE<sub>2</sub> concentrations  $\geq 30$  nM, progesterone production increased progressively by up to  $91 \pm 6\%$  (at 3000 nM PGE<sub>2</sub>) relative to the control rate of progesterone output in the absence of PGE<sub>2</sub> ( $P < 0.01$ ). However, co-treatment with SC19220 attenuated the responses to both 1000 and 3000 nM PGE<sub>2</sub> by  $56 \pm 4\%$  ( $P < 0.001$ ) and  $35 \pm 3\%$  ( $P < 0.001$ ) respectively, while co-treatment with AH6809 completely

**Table 1** Effects of SC19220 and AH6809 on basal progesterone production and 11 $\beta$ -hydroxysteroid dehydrogenase (11 $\beta$ HSD1) activity. Values are the mean  $\pm$  S.E.M. for three independent experiments with quadruplicate determinations in each experiment (one-way ANOVA)

	Progesterone (pmol/10 <sup>5</sup> cells)	11 $\beta$ HSD1 activity (pmol cortisone/4 h .5 $\times 10^4$ cells)
Control	2351.1 $\pm$ 1028.3	3.7 $\pm$ 1.2
+SC19220	2041.6 $\pm$ 913.6	3.8 $\pm$ 1.1
+AH6809	2712.5 $\pm$ 1318.1	2.5 $\pm$ 0.4
ANOVA	$P = 0.161$	$P = 0.158$

Progesterone concentrations in culture medium and 11 $\beta$ HSD1 activities in human granulosa–lutein cells were each assessed following a 24- or 4-h incubation respectively with 10  $\mu$ M SC19220 or 10  $\mu$ M AH6809.




**Figure 1** Effects of SC19220 and AH6809 on the progesterone responses to PGE<sub>2</sub> by human granulosa–lutein cells. Progesterone concentrations in medium collected after a 24-h incubation in the presence of 0–3000 nM PGE<sub>2</sub>, either in the absence (circle) or in the presence of 10 µM SC19220 (square) or 10 µM AH6809 (triangle). Values are the mean  $\pm$  S.E.M. for three independent experiments with quadruplicate determinations in each experiment. \* $P$ <0.05, \*\* $P$ <0.01, \*\*\* $P$ <0.001 relative to 0 nM PGE<sub>2</sub> in the absence or presence of the PTGER receptor antagonist, as appropriate. <sup>a</sup> $P$ <0.05, <sup>c</sup> $P$ <0.001 relative to cells treated with the same PGE<sub>2</sub> concentration in the absence of PTGER receptor antagonists (one-way ANOVA plus Bonferroni's *post hoc* multiple comparison test).

abolished the ability of all concentrations of PGE<sub>2</sub> to stimulate progesterone output.

Incubation for 24 h with PGE<sub>2</sub> evoked a concentration-dependent increase in cAMP accumulation which only achieved significance at the supraphysiological concentrations of 1000 and 3000 nM PGE<sub>2</sub> (Fig. 2). At the highest tested concentration of 3000 nM, PGE<sub>2</sub> increased cAMP accumulation by  $18.7 \pm 6.8$ -fold ( $P$ <0.01) in the absence of PTGER receptor antagonists. In cells co-treated with SC19220, this maximal concentration of PGE<sub>2</sub> increased cAMP accumulation by  $40.3 \pm 27.8$ -fold ( $P$ <0.01), whereas co-treatment with AH6809 suppressed the cAMP response to 3000 nM PGE<sub>2</sub> to just  $4.1 \pm 1.2$ -fold ( $P$ <0.05; Fig. 2).

#### Effects of SC19220 and AH6809 on 11 $\beta$ HSD1 activities in response to PGE<sub>2</sub> and butaprost in human granulosa–lutein cells

Consistent with our recent report (Jonas *et al.* 2006), PGE<sub>2</sub> significantly increased 11 $\beta$ HSD1 activity in a concentration-dependent manner; net oxidation of cortisol to cortisone was increased by up to  $43 \pm 3\%$  at the highest tested PGE<sub>2</sub> concentration of 1000 nM ( $P$ <0.001; Fig. 3). Co-treatment with SC19220 further enhanced 11 $\beta$ HSD1 activities at each of the tested concentrations of PGE<sub>2</sub>. At the highest tested concentration of 1000 nM PGE<sub>2</sub>, co-treatment with SC19220 increased the 11 $\beta$ HSD1 activity by a further  $32.7 \pm 10.2\%$  ( $P$ <0.05) relative to the cells treated with PGE<sub>2</sub> in the absence



**Figure 2** Effects of SC19220 and AH6809 on the cAMP responses to PGE<sub>2</sub> by human granulosa–lutein cells. Concentrations of cAMP after a 24-h incubation in the presence of 0–3000 nM PGE<sub>2</sub>, either in the absence (circle) or in the presence of 10 µM SC19220 (square) or 10 µM AH6809 (triangle). Values are the mean  $\pm$  S.E.M. for three independent experiments with quadruplicate determinations in each experiment. \* $P$ <0.05, \*\* $P$ <0.01 relative to 0 nM PGE<sub>2</sub> in the absence or presence of the PTGER receptor antagonists, as appropriate (one-way ANOVA plus Bonferroni's *post hoc* multiple comparison test).

of PTGER receptor antagonists. In contrast, co-treatment with AH6809 completely abolished the 11 $\beta$ HSD1 response to all tested concentrations of PGE<sub>2</sub> (Fig. 3).


Incubation for 4 h with the preferential PTGER2 receptor agonist butaprost also elicited a concentration-dependent increase in 11 $\beta$ HSD1 activities (Fig. 4). At the highest tested concentration of 1000 nM, butaprost increased net cortisol oxidation by  $40 \pm 3\%$  ( $P$ <0.01). Co-treatment with SC19220 further increased 11 $\beta$ HSD1 activity at each tested concentration of butaprost by up to  $31 \pm 1\%$  ( $P$ <0.05) relative to cells treated with butaprost alone. In contrast, as for PGE<sub>2</sub>, co-treatment with AH6809 completely abolished the stimulation of 11 $\beta$ HSD1 activities by each concentration of butaprost.

## Discussion


This study has investigated the potential participation of PTGER1 and PTGER2 receptors in mediating the effects of PGE<sub>2</sub> on progesterone production and glucocorticoid metabolism in human granulosa–lutein cells. Using preferential pharmacological agonists and antagonists for the PTGER1 and PTGER2 receptor subtypes, the combined data implicate PTGER2 receptors as mediators of the stimulation of progesterone synthesis and 11 $\beta$ HSD1 activity by PGE<sub>2</sub> in this human ovarian cell model *in vitro*.

The ability of PGE<sub>2</sub> to stimulate progesterone synthesis in human and non-human primate luteal cells, and in the widely used model of human granulosa–lutein cells, is well documented (Hahlin *et al.* 1988, Fehrenbach *et al.* 1999,





**Figure 3** Effects of PGE<sub>2</sub> on 11βHSD1 activities human granulosa-lutein cells in the absence and presence of SC19220 and AH6809. 11βHSD1 activities following treatment of cells with 0–1000 nM PGE<sub>2</sub> either in the absence (circle) or in the presence of 10 μM SC19220 (square) or 10 μM AH6809 (triangle). Values are the mean ± S.E.M. for three independent experiments with triplicate determinations in each experiment. \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001 relative to 0 nM PGE<sub>2</sub> in the absence or presence of the PTGER receptor antagonists, as appropriate. <sup>a</sup>*P* < 0.05, <sup>b</sup>*P* < 0.01 relative to cells treated with the same PGE<sub>2</sub> concentration in the absence of PTGER receptor antagonists (one-way ANOVA plus Bonferroni's *post hoc* multiple comparison test).



**Figure 4** Effects of butaprost on 11βHSD1 activities in human granulosa-lutein cells in the absence and presence of SC19220 and AH6809. 11βHSD1 activities following treatment of cells with 0–1000 nM butaprost either in the absence (circle) or in the presence of 10 μM SC19220 (square) or 10 μM AH6809 (triangle). Values are the mean ± S.E.M. for three independent experiments with triplicate determinations in each experiment. \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001 relative to 0 nM butaprost in the absence or presence of the PTGER receptor antagonists, as appropriate. <sup>a</sup>*P* < 0.05, <sup>b</sup>*P* < 0.01 relative to cells treated with the same butaprost concentration in the absence of PTGER receptor antagonists (one-way ANOVA plus Bonferroni's *post hoc* multiple comparison test).

Vaananen *et al.* 2001). Given the reported ability of PGE<sub>2</sub> to increase intracellular cAMP concentrations in these cells (Michael *et al.* 1993) and the established role for cAMP as a second messenger that stimulates steroidogenesis (Cooke 1999), it has widely been assumed that the progesterone response to PGE<sub>2</sub> is mediated via this cyclic nucleotide. We have previously demonstrated that both progesterone production and cAMP accumulation in human granulosa-lutein cells can be stimulated by the mixed PTGER1/PTGER2 receptor agonist, dimethyl-PGE<sub>2</sub> and by the preferential PTGER2 receptor agonist, butaprost (Harris *et al.* 2001). The cAMP responses to dimethyl-PGE<sub>2</sub> and butaprost were completely abolished by co-treatment with the PTGER1/PTGER2 receptor antagonist AH6809, but not the selective PTGER1 receptor antagonist SC19220, implicating the PTGER2 receptor in the cAMP response to these pharmacological agonists (Harris *et al.* 2001). Interestingly, the cAMP response to PGE<sub>2</sub> was only partially suppressed (by around 60%) with AH6809, and we did not assess the ability of these compounds to interfere with the steroidogenic responses to PGE<sub>2</sub> or its functional analogues.

In the present study, when cells were co-treated with SC19220, the progesterone response to PGE<sub>2</sub> was partially inhibited but not completely abolished. This suggests that the PTGER1 receptors participate in the stimulation of steroidogenesis by PGE<sub>2</sub>, but do not mediate the full steroidogenic response. We have now confirmed that co-treatment with AH6809 could completely prevent the steroidogenic response to PGE<sub>2</sub> at all tested concentrations, implicating an AH6809-sensitive receptor in the steroidogenic response to PGE<sub>2</sub>.

The concentrations of PGE<sub>2</sub> (and butaprost) used for the studies reported herein were selected based on previous investigations of human granulosa-lutein cells, which routinely use PGs at supraphysiological concentrations of 500–5000 nM. It is noteworthy that, at the lower tested concentrations of 3 and 10 nM, PGE<sub>2</sub> failed to stimulate steroidogenesis, even though Abramovitz *et al.* (2000) have previously reported that PGE<sub>2</sub> concentrations of < 30 nM should be sufficient to stimulate all four PTGER receptor subtypes. Moreover, although PGE<sub>2</sub> and butaprost were able to stimulate progesterone synthesis (and cortisol metabolism) at concentrations of 30 nM and above, at these concentrations, PGE<sub>2</sub> would be expected to saturate all four classes of PTGER receptor and possibly to activate non-PTGER receptors (i.e. PTGFR and D series prostaglandin (PTGDR) receptors), particularly at the conventional concentrations of 300, 1000 and 3000 nM (Abramovitz *et al.* 2000). Since PTGFR receptors are known to inhibit (rather than stimulate) progesterone production (Sugimoto *et al.* 1997, Tsai *et al.* 1998, 2001), it seems highly unlikely that the steroidogenic response to PGE<sub>2</sub> (even at concentrations > 30 nM) is mediated via the PTGFR receptor. However, we cannot yet exclude participation of the PTGDR receptor in the stimulation of steroidogenesis at high concentrations of PGE<sub>2</sub>, given that AH6809 has also been reported to function

as a weak PTGDR antagonist when used at a concentration of 10  $\mu$ M (Keery & Lumley 1988, Woodward *et al.* 1995).

In terms of effects on cAMP accumulation, we found that although PGE<sub>2</sub> increased cAMP accumulation in a concentration-dependent manner, this response only achieved statistical significance at the higher concentrations of 1000 and 3000 nM PGE<sub>2</sub>, which would be expected to bind all PTGER maximally, and possibly to activate PTGDR and PTGFR receptors as well (Abramovitz *et al.* 2000). Co-treatment of cells with AH6809 suppressed the PGE<sub>2</sub>-induced increase in the cAMP accumulation, but did not completely abolish the cAMP response to PGE<sub>2</sub>, indicating that the residual effect of high PGE<sub>2</sub> concentrations on cAMP may be mediated through an AH6809-insensitive receptor. SC19220, on the other hand, did not suppress the responsiveness of the cells to PGE<sub>2</sub>. If anything, co-treatment with the PTGER1 antagonist SC19220 enhanced the increase of cAMP in response to PGE<sub>2</sub>. These observations raise two possibilities. First, inhibition of PTGER1 receptor binding by SC19220 may have facilitated increased binding of PGE<sub>2</sub> to other prostanoid receptors which are capable of stimulating cAMP accumulation (e.g. PTGER2, PTGER4 and/or PTGDR receptors). Secondly, co-stimulation of the PTGER1 receptor would be expected to activate the calcium-protein kinase C signalling pathway (Narumiya *et al.* 1999), which can inhibit the stimulation of cAMP production (Abayasekara *et al.* 1993a,b). Based on these observations, prostanoid receptors which are sensitive to AH6809 but insensitive to SC19220 (e.g. PTGER2 and/or PTGDR) are implicated in the ability of PGE<sub>2</sub> to stimulate cAMP accumulation in human granulosa-lutein cells. In light of our previous report (Harris *et al.* 2001), we would suggest that while the ability of dimethyl-PGE<sub>2</sub> and butaprost to elevate intracellular cAMP concentrations is mediated solely via a PG receptor that can be antagonised with AH6809, consistent with a role for the PTGER2 receptor in eliciting a cAMP response, at least some of the second messenger response to PGE<sub>2</sub> is mediated via AH6809-resistant receptors, which may include PTGER4 receptors and possibly even PTGDR receptors. We also note that, in the present study, the residual cAMP responses to the higher concentrations of PGE<sub>2</sub> in cells co-treated with AH6809 were not accompanied by any significant increases in progesterone synthesis, showing that second messengers other than cAMP may be required for PGE<sub>2</sub> to stimulate steroidogenesis.

Turning to the regulation of cortisol-cortisone metabolism by PGE<sub>2</sub>, we have previously established that luteinizing human granulosa cells only express 11 $\beta$ HSD1 with no detectable expression of 11 $\beta$ HSD2 protein or mRNA, even when the PCR is used to increase the sensitivity of detection (Michael *et al.* 1997, Thurston *et al.* 2003a). Although the 11 $\beta$ HSD1 enzyme is intrinsically bidirectional, in most tissues this enzyme acts predominantly as an 11-ketosteroid reductase to regenerate cortisol from cortisone (Seckl & Walker 2001, Seckl 2004, Draper & Stewart 2005). However, we have previously found that 11 $\beta$ HSD1 acts predominantly, if not

exclusively, to catalyse the oxidative inactivation of cortisol in human granulosa-lutein cells (Michael *et al.* 1997, Thurston *et al.* 2003a), bovine granulosa and luteal cells (Thurston *et al.* 2007) and porcine granulosa cells (Sunak *et al.* 2007). 11 $\beta$ HSD1 has also been reported to act predominantly as a 11 $\beta$ -dehydrogenase in rat testis Leydig cells (Gao *et al.* 1997, Ge & Hardy 2000) suggesting that the major action of the 11 $\beta$ HSD1 enzyme may be fundamentally different in the steroidogenic cells of the ovary and testis (Michael *et al.* 2003, Ge *et al.* 2005). Recent studies have established that the predominant direction of 11 $\beta$ HSD1 is dependent on the ratio of the reduced form of nicotinamide adenine dinucleotide phosphate to its oxidised form (NADPH: NADP<sup>+</sup>) within the lumen of the smooth endoplasmic reticulum (Draper *et al.* 2003, Atanasov *et al.* 2004, Banhegyi *et al.* 2004, Bujalska *et al.* 2005, McCormick *et al.* 2006) such that changes in this ratio in steroidogenic gonadal cells may favour the oxidative activity of 11 $\beta$ HSD1 in these cells (Michael *et al.* 2003, Ge *et al.* 2005).

In the present study, the stimulation of the net oxidation of cortisol in human granulosa-lutein cells by PGE<sub>2</sub> showed good agreement with our recently published data (Jonas *et al.* 2006). The fact that this stimulation could be reproduced using the preferential PTGER2 receptor agonist butaprost, combined with the fact that AH6809 could completely suppress the effects of PGE<sub>2</sub> and butaprost on cortisol oxidation, suggests that the stimulation of the oxidative activity of 11 $\beta$ HSD1 is mediated via the PTGER2 receptor. In contrast, co-treatment with SC19220 enhanced the 11 $\beta$ HSD1 response at each of the tested concentrations of PGE<sub>2</sub>, suggesting that the co-activation of PTGER1 receptors limits the stimulation of cortisol-cortisone metabolism by PGE<sub>2</sub>. PTGER1 receptors are known to act via calcium as a second messenger, and we have confirmed that in human granulosa-lutein cells, SC19220 can antagonise the ability of PGE<sub>2</sub> and its dimethyl derivative to increase the intracellular calcium concentration (Harris *et al.* 2001). Given that studies of placental biochemistry have shown that calcium can mediate the inhibition of 11 $\beta$ HSD1 activity in response to PGs and leukotriene B<sub>4</sub> (Hardy *et al.* 1999, 2001), we would propose that activation of the PTGER1 receptor-calcium signalling pathway by PGE<sub>2</sub> limits the ability of this PG to stimulate 11 $\beta$ HSD1 in human granulosa-lutein cells via the PTGER2-cAMP pathway.

In comparing the three responses considered in this study, we note that at the lower concentrations tested (<1000 nM), which are likely to fall within the physiological range and selectively to activate only the PTGER receptor subtypes, PGE<sub>2</sub> can stimulate both progesterone synthesis and cortisol metabolism without exerting a significant effect on the intracellular cAMP concentration. This suggests that either the effects of 10–300 nM PGE<sub>2</sub> on steroid synthesis and metabolism are mediated via local increases in cAMP within specific subcellular microenvironments (not reflected by changes in the total intracellular cAMP concentration) or

these effects are mediated via a second messenger other than cAMP.

In summary, we have presented data which implicate PTGER2 receptors in the stimulation of progesterone synthesis, cAMP accumulation and cortisol metabolism via 11 $\beta$ HSD by PGE<sub>2</sub> in human granulosa–lutein cells. Further studies are necessary to elucidate potential roles for other prostanoid receptors, such as PTGDR receptors, in the responses at the upper concentrations of PGE<sub>2</sub>.

## Acknowledgements

We wish to thank the staff at the Lister Private Hospital assisted conception unit (Chelsea, London, UK) for providing the human granulosa–lutein cells used for this study. This work was supported by a Wellcome Trust Grant No. 056630. For this publication, the contemporary HUGO nomenclature has been used for the prostanoid receptors, where PTGER1, PTGER2, PTGER3, PTGER4, PTGDR and PTGFR receptors have been previously described as EP<sub>1</sub>, EP<sub>2</sub>, EP<sub>3</sub>, EP<sub>4</sub>, DP and FP receptors respectively. According to the HUGO nomenclature, the hydroxysteroid dehydrogenase enzymes type 1 11 $\beta$ HSD/11 $\beta$ HSD1 and type 2 11 $\beta$ HSD/11 $\beta$ HSD2 have been designated as HSD11B1 and HSD11B2 respectively; however, for this publication, the conventional (pre-HUGO) nomenclature has been used for the 11 $\beta$ HSD enzymes. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

## References

- Abayasekara DRE, Jones PM, Persaud SJ, Michael AE & Flint APF 1993a Prostaglandin F<sub>2 $\alpha$</sub>  activates protein kinase C in human ovarian cells. *Molecular and Cellular Endocrinology* **91** 51–57.
- Abayasekara DRE, Michael AE, Webley GE & Flint APF 1993b Mode of action of prostaglandin F<sub>2 $\alpha$</sub>  in human luteinized granulosa cells: role of protein kinase C. *Molecular and Cellular Endocrinology* **97** 81–91.
- Abramovitz M, Adam M, Boie Y, Carriere MC, Denis D, Godbout C, Lamontagne S, Rochette C, Sawyer N, Tremblay NM *et al.* 2000 The utilisation of recombinant prostanoid receptors to determine the affinities and selectivities of prostaglandins and related analogues. *Biochimica et Biophysica Acta* **1483** 285–293.
- Alfaidy N, Xiong ZG, Myatt L, Lye SJ, Macdonald JF & Challis JR 2001 Prostaglandin F<sub>2 $\alpha$</sub>  potentiates cortisol production by stimulating 11 $\beta$ -hydroxysteroid dehydrogenase 1: a novel feedback loop that may contribute to human labor. *Journal of Clinical Endocrinology and Metabolism* **86** 5585–5592.
- Atanasov AG, Nashev LG, Schweizer RAS, Frick C & Odermatt A 2004 Hexose-6-phosphate dehydrogenase determines the reaction direction of 11 $\beta$ -hydroxysteroid dehydrogenases type 1 as an oxoreductase. *FEBS Letters* **571** 129–133.
- Banhegyi G, Benedetti A, Fulceri R & Senesi S 2004 Co-operativity between 11 $\beta$ -hydroxysteroid dehydrogenases type 1 and hexose-6-phosphate dehydrogenase in the lumen of the endoplasmic reticulum. *Journal of Biological Chemistry* **279** 27017–27021.
- Bujalska IJ, Draper N, Michailidou Z, Tomlinson JW, White PC, Chapman KE, Walker EA & Stewart PM 2005 Hexose-6-phosphate dehydrogenase confers oxo-reductase activity upon 11 $\beta$ -hydroxysteroid dehydrogenase type 1. *Journal of Molecular Endocrinology* **34** 675–684.
- Challis JR, Lye SJ & Gibb W 1997 Prostaglandins and parturition. *Annals of the New York Academy of Science* **26** 254–267.
- Challis JR, Matthews SG, Gibb W & Lye SJ 2000 Endocrine and paracrine regulation of birth at term and preterm. *Endocrine Reviews* **21** 514–550.
- Chen WY, Yang JG & Li PS 2000 Effect of dexamethasone on the expression of p34(cdc2) and cyclin B1 in pig oocytes *in vitro*. *Molecular Reproduction and Development* **56** 74–79.
- Coleman RA, Smith WL & Narumiya S 1994 International union of pharmacology classification of prostanoid receptors: properties, distribution, and structure of the receptors and their subtypes. *Pharmacological Reviews* **46** 205–229.
- Cooke BA 1999 Signal transduction involving cyclic AMP-dependent and cyclic AMP-independent mechanisms in the control of steroidogenesis. *Molecular and Cellular Endocrinology* **151** 25–35.
- Draper N & Stewart PM 2005 11 $\beta$ -hydroxysteroid dehydrogenase and the pre-receptor regulation of corticosteroid hormone action. *Journal of Endocrinology* **186** 251–271.
- Draper N, Walker EA, Bujalska IJ, Tomlinson JW, Chalder SM, Arlt W, Lavery GG, Bedendo O, Ray DW, Laing I *et al.* 2003 Mutations in the genes encoding 11 $\beta$ -hydroxysteroid dehydrogenase type 1 and hexose-6-phosphate dehydrogenase interact to cause cortisone reductase deficiency. *Nature Genetics* **34** 434–439.
- Fehrenbach A, Hodges JK & Einspanier A 1999 Direct effects of the prostaglandins E<sub>2</sub> and F<sub>2 $\alpha$</sub>  on progesterone release by the corpus luteum of the marmoset monkey (*Callithrix jacchus*) studied by *in vitro* microdialysis. *Journal of Endocrinology* **161** 433–443.
- Fowkes RC, Chandras C, Chin EC, Okolo S, Abayasekara DRE & Michael AE 2001 Relationship between the production of prostaglandins and progesterone by luteinizing human granulosa cells. *Journal of Endocrinology* **171** 455–462.
- Funk CD, Furci L, FitzGerald GA, Grugorzcyk R, Rochette C, Bayne MA, Abramovitz M, Adam M & Metters KM 1993 Cloning and expression of a cDNA for the human prostaglandin E receptor EP1 subtype. *Journal of Biological Chemistry* **268** 26767–26772.
- Gao HB, Ge RS, Lakshmi V, Marandici A & Hardy MP 1997 Hormonal regulation of oxidative and reductive activities of 11 $\beta$ -hydroxysteroid dehydrogenase in rat Leydig cells. *Endocrinology* **138** 156–161.
- Ge RS & Hardy MP 2000 Initial predominance of the oxidative activity of type I 11 $\beta$ -hydroxysteroid dehydrogenase in primary rat Leydig cells and transfected cell lines. *Journal of Andrology* **21** 303–310.
- Ge RS, Dong Q, Niu EM, Sottas CM, Hardy DO, Catterall JF, Latif SA, Morris DJ & Hardy MP 2005 11 $\beta$ -hydroxysteroid dehydrogenase 2 in rat Leydig cells: its role in blunting glucocorticoid action at physiological levels of substrate. *Endocrinology* **146** 2657–2664.
- Greeley MS, Calder DR, Taylor MH, Hols H & Wallace RA 1986 Oocyte maturation in mummichog (*Fundulus heteroclitus*): effects of steroids on germinal vesicle breakdown of intact follicles. *General and Comparative Endocrinology* **62** 281–289.
- Hahlin M, Dennefors B, Johanson C & Hamberger L 1988 Luteotropic effects of prostaglandin E<sub>2</sub> on the human corpus luteum of the menstrual cycle and early pregnancy. *Journal of Clinical Endocrinology and Metabolism* **66** 909–914.
- Hardy DB, Pereria LE & Yang K 1999 Prostaglandins and leukotriene B<sub>4</sub> are potent inhibitors of 11 $\beta$ -hydroxysteroid dehydrogenase type 2 activity in human choriocarcinoma JEG-3 cells. *Biology of Reproduction* **61** 40–45.
- Hardy DB, Dixon SJ, Narayanan N & Yang K 2001 Calcium inhibits human placental 11 $\beta$ -hydroxysteroid dehydrogenase type 2 activity. *Biochemical and Biophysical Research Communications* **283** 756–761.
- Harlow CR, Jenkins JM & Winston RML 1997 Increased follicular fluid total and free cortisol levels during the luteinizing hormone surge. *Fertility and Sterility* **68** 48–53.
- Harper JF & Brooker G 1975 Femtomole sensitive radioimmunoassay for cyclic AMP and cyclic GMP after 2'0 acetylation by acetic anhydride in aqueous solution. *Journal of Cyclic Nucleotide Research* **1** 207–218.

- Harris TE, Squires PE, Michael AE, Lopez-Bernal A & Abayasekara DRE 2001 Human granulosa-lutein cells express functional EP<sub>1</sub> and EP<sub>2</sub> prostaglandin receptors. *Biochemical and Biophysical Research Communications* **285** 1089–1094.
- Hirata M, Hayashi Y, Ushikubi F, Yokota Y, Kageyama R, Nakanishi S & Narumiya S 1991 Cloning and expression of cDNA for a human thromboxane A<sub>2</sub> receptor. *Nature* **349** 617–620.
- Honda A, Sugimoto Y, Namba T, Watabe A, Irie A, Negishi M, Narumiya S & Ichikawa A 1993 Cloning and expression of a cDNA for mouse prostaglandin E receptor EP<sub>2</sub> subtype. *Journal of Biological Chemistry* **268** 7759–7762.
- Jonas KC, Chandras C, Abayasekara DRE & Michael AE 2006 Role for prostaglandins in the regulation of type 1 11 $\beta$ -hydroxysteroid dehydrogenase in human granulosa-lutein cells. *Endocrinology* **147** 5865–5872.
- Keery RJ & Lumley P 1988 AH6809, a prostaglandin DP-receptor blocking drug on human platelets. *British Journal of Pharmacology* **94** 745–754.
- López Bernal A, Anderson ABM & Turnbull AC 1982a Cortisol:cortisone interconversion by human decidua in relation to parturition. Effect of tissue manipulation on 11 $\beta$ -hydroxysteroid dehydrogenase activity. *Journal of Endocrinology* **93** 141–149.
- López Bernal A, Anderson ABM & Turnbull AC 1982b The lack of influence of parturition on human placental 11 $\beta$ -hydroxysteroid dehydrogenase activity. *Journal of Clinical Endocrinology and Metabolism* **54** 1251–1254.
- McCormick KL, Wang X & Mick GJ 2006 Evidence that the 11 $\beta$ -hydroxysteroid dehydrogenase (11 $\beta$ HSD1) is regulated by pentose pathway flux: studies in rat adipocytes and microsomes. *Journal of Biological Chemistry* **281** 341–347.
- Michael AE, Abayasekara DRE & Webley GE 1993 The luteotrophic actions of prostaglandin E<sub>2</sub> and F<sub>2 $\alpha$</sub>  are differentially mediated via cyclic AMP and protein kinase C. *Journal of Endocrinology* **138** 291–298.
- Michael AE, Abayasekara DRE & Webley GE 1994 Cellular mechanisms of luteolysis. *Molecular and Cellular Endocrinology* **99** R1–R9.
- Michael AE, Gregory L, Piercy EC, Walker SM, Shaw RW & Cooke BA 1995 Ovarian 11 $\beta$ -hydroxysteroid dehydrogenase activity is inversely related to the outcome of in vitro fertilization-embryo transfer treatment cycles. *Fertility and Sterility* **64** 590–598.
- Michael AE, Evangelatou M, Norgate DP, Clarke RJ, Antoniwi JW, Stedman B, Brennan A, Welsby R, Bujalska I & Stewart PM 1997 Isoforms of 11 $\beta$ -hydroxysteroid dehydrogenase in human granulosa-lutein cells. *Molecular and Cellular Endocrinology* **132** 43–52.
- Michael AE, Thurston LM & Rae MT 2003 Glucocorticoid metabolism and reproduction: a tale of two enzymes. *Reproduction* **126** 425–441.
- Narko K, Saukkonen K, Ketola I, Butzow R, Heikinheimo M & Ristimäki A 2001 Regulated expression of prostaglandin E<sub>2</sub> receptors EP<sub>2</sub> and EP<sub>4</sub> in human ovarian granulosa cells. *Journal of Clinical Endocrinology and Metabolism* **86** 1765–1768.
- Narumiya S, Sugimoto Y & Ushikubi F 1999 Prostanoid receptors: structures, properties and functions. *Physiological Reviews* **79** 1193–1226.
- Olofsson J & Leung PCK 1994 Autocrine/paracrine role of prostaglandins in corpus luteum function. *Molecular and Cellular Endocrinology* **100** 87–91.
- Pallikaros Z, Schulster D, Baldwin SA, Helliwell RJA, Michael AE & Cooke BA 1995 Characterization of site-directed antibodies to the LH receptor in functionally active gonadal cells and their differential effects on LH-stimulated signal transduction in Leydig tumor (MA-10) cells. *Molecular and Cellular Endocrinology* **114** 57–68.
- Richardson MC 1986 Hormonal control of ovarian luteal cells. *Oxford Reviews in Reproductive Biology* **8** 321–378.
- Seckl JR & Walker BR 2001 Mini-review: 11 $\beta$ -hydroxysteroid dehydrogenase type 1 – a tissue-specific amplifier of glucocorticoid action. *Endocrinology* **142** 1371–1376.
- Seckl JR 2004 11 $\beta$ -hydroxysteroid dehydrogenases: changing glucocorticoid action. *Current Opinion in Pharmacology* **4** 597–602.
- Steiner AL, Wehmann RE, Parker CW & Kipnis DM 1972 Radioimmunoassay for the measurement of cyclic nucleotides. *Advances in Cyclic Nucleotide Research* **2** 51–61.
- Sugimoto Y, Namba T, Shigemoto R, Negishi M, Ichikawa A & Narumiya S 1992 Cloning and expression of a cDNA for mouse prostaglandin E receptor EP<sub>3</sub> subtype. *Journal of Biological Chemistry* **267** 6463–6466.
- Sugimoto Y, Yamasaki A, Segi E, Tsuboi K, Aze Y, Nishimura T, Oida H, Yoshida N, Tanaka T, Katsuyama M *et al.* 1997 Failure of parturition in mice lacking the prostaglandin F receptor. *Science* **277** 681–683.
- Sunak N, Green DF, Abeydeera LR, Thurston LM & Michael AE 2007 Implication of cortisol and 11 $\beta$ -hydroxysteroid dehydrogenase enzymes in the development of porcine ovarian follicles and cysts. *Reproduction*, **133** 1149–1158.
- Tetsuka M, Thomas FJ, Thomas MJ, Anderson RA, Mason JI & Hillier SG 1997 Differential expression of messenger ribonucleic acids encoding 11 $\beta$ -hydroxysteroid dehydrogenase types 1 and 2 in human granulosa cells. *Journal of Clinical Endocrinology and Metabolism* **82** 2006–2009.
- Tetsuka M, Haines LC, Milne M, Simpson GE & Hillier SG 1999 Regulation of 11 $\beta$ -hydroxysteroid dehydrogenase type 1 gene expression by LH and interleukin-1 $\beta$  in cultured rat granulosa cells. *Journal of Endocrinology* **163** 417–423.
- Thurston LM, Chin E, Jonas KC, Bujalska IJ, Stewart PM, Abayasekara DRE & Michael AE 2003a Expression of 11 $\beta$ -hydroxysteroid dehydrogenase (11 $\beta$ HSD) proteins in luteinizing human granulosa-lutein cells. *Journal of Endocrinology* **178** 127–135.
- Thurston LM, Norgate DP, Jonas KC, Gregory L, Wood PJ, Cooke BA & Michael AE 2003b Ovarian modulators of type 1 11 $\beta$ -hydroxysteroid dehydrogenase (11 $\beta$ HSD) activity and intra-follicular cortisol:cortisone ratios correlate with the clinical outcome of IVF. *Human Reproduction* **18** 1603–1612.
- Thurston LM, Abayasekara DRE & Michael AE 2007 11 $\beta$ -Hydroxysteroid dehydrogenase expression and activities in bovine granulosa cells and corpora lutea implicate corticosteroids in bovine ovarian physiology. *Journal of Endocrinology* **193** 299–310.
- Tsai SJ & Wiltbank MC 1998 Prostaglandin F<sub>2 $\alpha$</sub>  regulates distinct physiological changes in early and mid-cycle bovine corpora lutea. *Biology of Reproduction* **58** 346–352.
- Tsai SJ, Wu MH, Chuang PC & Chen HM 2001 Distinct regulation of gene expression by prostaglandin F<sub>2 $\alpha$</sub>  (PGF<sub>2 $\alpha$</sub> ) is associated with PGF<sub>2 $\alpha$</sub>  resistance or susceptibility in human granulosa-luteal cells. *Molecular Human Reproduction* **7** 415–423.
- Vaananen JE, Tong BL, Vaananen CC, Chan IH, Yuen BH & Leung PC 2001 Interaction of prostaglandin F<sub>2 $\alpha$</sub>  and prostaglandin E<sub>2</sub> on progesterone production in human granulosa-luteal cells. *Biological Signals and Receptors* **10** 380–388.
- Watabe A, Sugimoto Y, Honda A, Irie A, Namba T, Negishi M, Ito S, Narumiya S & Ichikawa A 1993 Cloning and expression of cDNA for a mouse EP<sub>1</sub> subtype of prostaglandin E receptor. *Journal of Biological Chemistry* **268** 20175–20178.
- Webley GE, Luck MR & Hearn JP 1988 Stimulation of progesterone secretion by cultured human granulosa cells with melatonin and catecholamines. *Journal of Reproduction and Fertility* **84** 669–677.
- Whittle WL, Patel FA, Alfaidy N, Holloway AC, Fraser M, Gyomai S, Lye SJ, Gibb W & JR 2001 Glucocorticoid regulation of human and ovine parturition: the relationship between fetal hypothalamic-pituitary-adrenal axis activation and intrauterine prostaglandin production. *Biology of Reproduction* **64** 1019–1032.
- Woodward DF, Pepper DJ, Burkey TH & Regan JW 1995 6-Isopropoxy-9-oxoxanthene-2-carboxylic acid (AH6809), a human EP<sub>2</sub> receptor antagonist. *Biochemical Pharmacology* **50** 1731–1733.
- Yang JG, Chen WY & Li PS 1999 Effects of glucocorticoids on maturation of pig oocytes and their subsequent fertilizing capacity *in vitro*. *Biology of Reproduction* **60** 929–936.

Received in final form 31 May 2007

Accepted 10 June 2007

Made available online as an Accepted Preprint  
13 June 2007